

学校编码: 10384

分类号_____密级_____

学号: 21720080150405

UDC_____

廈門大學

博 士 学 位 论 文

多氯联苯对小鼠附睾的毒性效应及机制的研究

Effects and mechanisms of polychlorinated biphenyls
on mouse epididymis

蔡嘉力

指导教师姓名: 陈奕欣 教授

专业名称: 遗传学

论文提交日期: 2011 年 11 月

论文答辩时间: 2011 年 11 月

学位授予日期: 2011 年 月

答辩委员会主席: _____

评 阅 人: _____

2011 年 11 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为()课题(组)的研究成果,获得()课题(组)经费或实验室的资助,在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

（ ） 1. 经厦门大学保密委员会审查核定的保密学位论文，
于 2013 年 12 月 1 日解密，解密后适用上述授权。

（ ☒ ） 2. 不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月 日

目 录

| | |
|----------------------------|----|
| 目录 | I |
| Contents | V |
| 摘要 | IX |
| ABSTRACT | XI |
| 第一部分 前言 | 1 |
| 1.1 多氯联苯的理化性质和污染现状 | 1 |
| 1.1.1 来源和理化性质 | 1 |
| 1.1.2 环境污染现状 | 3 |
| 1.2 多氯联苯的雄性生殖毒性效应与机制研究 | 6 |
| 1.2.1 多氯联苯的生物毒性效应 | 6 |
| 1.2.2 PCB 对雄性生殖的影响 | 8 |
| 1.2.3 PCB 对雄性生殖系统毒性机制的研究进展 | 10 |
| 1.3 精子在附睾中的成熟以及污染物对附睾功能的影响 | 13 |
| 1.3.1 附睾的结构与功能 | 13 |
| 1.3.2 附睾的微环境与精子的成熟 | 14 |
| 1.3.3 附睾功能的调控机制 | 17 |
| 1.3.4 毒物对精子附睾成熟过程的影响 | 18 |
| 1.4 毒理基因组学的研究概况 | 20 |
| 1.4.1 毒理基因组学的概念 | 20 |
| 1.4.2 毒理基因组学在雄性生殖毒性研究中的应用 | 21 |
| 1.5 本研究的目的意义和主要内容 | 22 |
| 1.5.1 本研究目的和意义 | 22 |
| 1.5.2 研究内容 | 23 |
| 第二部分 材料和方法 | 24 |

| | |
|--|-----------|
| 2.1 实验动物 | 24 |
| 2.2 仪器与试剂 | 24 |
| 2.2.1 主要仪器 | 24 |
| 2.2.2 主要试剂 | 25 |
| 2.3 实验动物处理 | 25 |
| 2.4 取样及样品处理 | 26 |
| 2.5 精子质量检测方法 | 26 |
| 2.5.1 试剂配制 | 26 |
| 2.5.2 游离尾睾的精子 | 27 |
| 2.5.3 精子活力的测定 | 27 |
| 2.5.4 精子存活率的测定 | 27 |
| 2.5.5 精子密度的测定 | 27 |
| 2.5.6 精子形态染色 | 28 |
| 2.6 组织学切片 | 28 |
| 2.6.1 H.E 染色 | 28 |
| 2.6.2 免疫组织化学 | 29 |
| 2.7 Western blot 分析 | 30 |
| 2.7.1 蛋白提取及含量测定 | 30 |
| 2.7.2 SDS-聚丙烯酰胺凝胶电泳 (SDS-PAGE) 与 Western blot 分析 | 30 |
| 2.8 生化指标的测定 | 32 |
| 2.8.1 血清激素水平的测定 | 32 |
| 2.8.2 精子顶体酶活性测定 | 32 |
| 2.9 小鼠肝脏中 PCB 含量的测定 | 33 |
| 2.9.1 样品预处理 | 33 |
| 2.9.2 气相色谱分析检测 | 34 |
| 2.10 基因表达谱分析 | 34 |
| 2.10.1 头睾组织总 RNA 的提取 | 34 |
| 2.10.2 对样品 RNA 进行荧光标记 | 35 |
| 2.10.3 杂交与清洗 | 35 |

| | |
|--|-----------|
| 2.10.4 芯片扫描与芯片图像的采集..... | 36 |
| 2.10.5 差异表达基因的生物信息学分析..... | 36 |
| 2.11 实时荧光定量 PCR (Real-time PCR) | 37 |
| 2.11.1 引物设计..... | 37 |
| 2.11.2 cDNA 模板的合成 | 38 |
| 2.11.3 实时荧光定量 PCR (Real-time PCR) | 38 |
| 2.12 Rho 蛋白活性的检测 | 39 |
| 2.13 紧密连接蛋白磷酸化水平的检测 | 40 |
| 2.13.1 试剂配制..... | 40 |
| 2.13.2 制备组织裂解物..... | 40 |
| 2.13.3 免疫共沉淀..... | 41 |
| 2.14 头睾上皮细胞顶端紧密连接透过性的硝酸镧示踪 | 41 |
| 2.14.1 试剂配制..... | 41 |
| 2.14.2 小鼠心脏灌注..... | 41 |
| 2.14.3 电镜观察..... | 42 |
| 2.15 数据统计与处理 | 42 |
| 第三部分 结果与分析 | 43 |
| 3.1 Aroclor 1254 对小鼠附睾的毒性效应 | 43 |
| 3.1.1 小鼠在 Aroclor 1254 暴露后 PCB 的机体负荷..... | 43 |
| 3.1.2 Aroclor 1254 对雄性小鼠精子指标的影响 | 44 |
| 3.1.3 Aroclor 1254 对小鼠附睾组织学形态的影响 | 46 |
| 3.1.4 Aroclor 1254 对小鼠附睾的内分泌干扰效应 | 49 |
| 3.2 Aroclor 1254 对小鼠附睾头睾基因表达的影响 | 51 |
| 3.2.1 不同剂量 Aroclor 1254 暴露对小鼠附睾差异表达基因的影响 | 51 |
| 3.2.2 差异表达基因的实时定量 PCR 验证 | 53 |
| 3.2.3 差异表达基因的功能注释..... | 54 |
| 3.3 Aroclor 1254 对小鼠附睾紧密连接及其调控机制的影响 | 61 |
| 3.3.1 Aroclor 1254 对小鼠附睾小 GTP 结合蛋白表达的影响..... | 61 |
| 3.3.2 Aroclor 1254 对小鼠附睾细胞骨架调节功能的影响 | 62 |

| | |
|---|------------|
| 3.3.3 Aroclor 1254 对小鼠附睾紧密连接的影响 | 64 |
| 3.4 PCB 暴露的潜在标志物的鉴定 | 68 |
| 3.4.1 差异表达基因中的潜在标志物..... | 68 |
| 3.4.2 Psap 表达与 Aroclor 1254 暴露的剂量效应关系..... | 69 |
| 3.4.3 Psap 表达与肝脏 Aroclor 1254 浓度以及精子指标的相关性..... | 71 |
| 3.4.4 Psap 表达与 Aroclor 1254 暴露的时间效应关系..... | 72 |
| 第四部分 讨论 | 73 |
| 4.1 Aroclor 1254 在小鼠肝脏中的富集 | 73 |
| 4.2 Aroclor 1254 暴露对体重和精子参数的影响 | 73 |
| 4.3 Aroclor 1254 暴露对雌激素和雄激素的干扰 | 74 |
| 4.4 Aroclor 1254 暴露对附睾头部表达谱的影响 | 76 |
| 4.4.1 Aroclor 1254 暴露附睾头部差异基因表达的影响 | 76 |
| 4.4.2 Aroclor 1254 暴露对基因表达模式影响的差异 | 77 |
| 4.4.3 不同剂量 Aroclor 1254 暴露对附睾头部效应机制的差异 | 78 |
| 4.5 Aroclor 1254 暴露对细胞骨架的影响 | 81 |
| 4.6 Aroclor 1254 暴露对紧密连接功能的影响 | 84 |
| 4.7 附睾 PCB 生殖毒性的效应标志物的筛选 | 88 |
| 第五部分 结论和展望 | 91 |
| 5.1 结论 | 91 |
| 5.2 本研究的创新点和贡献 | 92 |
| 5.3 展望 | 92 |
| 参考文献 | 94 |
| 附录 | 115 |
| 致谢 | 131 |

Contents

| | |
|--|-----------|
| Contents (in Chinese) | I |
| Contents (in English) | V |
| ABSTRACT (in Chinese) | IX |
| ABSTRACT (in English)..... | XI |
| Chapter 1 Introduction | 1 |
| 1.1 Phisical-chemical properties and pollution status of PCBs..... | 1 |
| 1.1.1 Source and phisical-chemical properties of PCBs | 1 |
| 1.1.2 Pollution status of PCBs | 3 |
| 1.2 Reseach of effects and mechanisms of PCBs on male reproduction | 6 |
| 1.2.1.Toxicological effects of PCBs | 6 |
| 1.2.2 Toxicological effects of PCBs on male reproduction..... | 8 |
| 1.2.3 Advances in the research of mechanisms of PCBs on on male reproduction..... | 10 |
| 1.3 Epididymal sperm maturation and the effects of envrionmental pollutants on epididymis | 13 |
| 1.3.1 Structure and function of epididynis..... | 13 |
| 1.3.2 Epididymal microenvironemnt and sperm maturation | 14 |
| 1.3.3 Regualtion of epididymis function..... | 17 |
| 1.3.4 Effects of toxicants on sperm maturation | 18 |
| 1.4 Application of toxicogenomics | 20 |
| 1.4.1 Concept of toxicogenomics..... | 20 |
| 1.4.2 Application of toxicogenomics in male reproductive toxicology..... | 21 |
| 1.5 Objectives and scientific significance of the study | 22 |
| Chapter 2 Materials and Methods..... | 24 |

| | |
|--|-----------|
| 2.1 Experimental speices | 24 |
| 2.2 Instruments and chemicals..... | 24 |
| 2.2.1 Major instruments | 24 |
| 2.2.2 Major chemicals..... | 25 |
| 2.3 Animal treatments..... | 25 |
| 2.4 Smapling and sample proceesing..... | 26 |
| 2.5 Determiantion of sperm parameters | 26 |
| 2.5.1 Reagent | 26 |
| 2.5.2 Isolation of epididymal sperm | 27 |
| 2.5.3 Determination of sperm motility..... | 27 |
| 2.5.4 Determination of sperm viability | 27 |
| 2.5.5 Determination of sperm concentration..... | 27 |
| 2.5.6 Mophological observation of sperm | 28 |
| 2.6 Histological examination | 28 |
| 2.6.1 Staining with H.E..... | 28 |
| 2.6.2 Immunohistochemistry | 29 |
| 2.7 Western blot assay..... | 30 |
| 2.7.1 Protein extraction and determination | 30 |
| 2.7.2 SDS-PAGE and Western blot analysis | 30 |
| 2.8 Biochemical analysis | 32 |
| 2.8.1 Determination of serum steroid hormones..... | 32 |
| 2.8.2 Determination acrosin activity | 32 |
| 2.9 Determination of hepatic PCB accumulation in mice | 33 |
| 2.9.1 Sample processing | 33 |
| 2.9.2 Gas chromatograph | 34 |
| 2.10 Gene expression profiles..... | 34 |
| 2.10.1 Isolation of total RNA of caput epididymis..... | 34 |
| 2.10.2 Fluorescence labeling the RNA sample..... | 35 |
| 2.10.3 Hybridization | 35 |
| 2.10.4 Chip scanning and image processing..... | 36 |

| | |
|--|-----------|
| 2.10.5 cDNA microarray data analysis | 36 |
| 2.11 Real-time fluorescent quantitative PCR | 37 |
| 2.11.1 Primers design..... | 37 |
| 2.11.2 Reverse transcription | 38 |
| 2.11.3 Real-time PCR | 38 |
| 2.12 Rho activity assay..... | 39 |
| 2.13 Determination of occludin phosphorylation status..... | 40 |
| 2.13.1 Reagent | 40 |
| 2.13.2 Preparation of tissue lyaste | 40 |
| 2.13.3 Immunoprecipitate | 41 |
| 2.14 Tracer studies on epididymla tight junctions | 41 |
| 2.14.1 Reagent | 41 |
| 2.14.2 Mice heart perfusion | 41 |
| 2.14.3 Electron microscope observation..... | 42 |
| 2.15 Statistical analysis | 42 |
| Chpater 3 Results and Analysis..... | 43 |
| 3.1 Effects of Aroclor 1254 on mouse epididymis..... | 43 |
| 3.1.1 PCB body burden in mice exposed to Aroclor 1254 | 43 |
| 3.1.2 Effects of Aroclor 1254 on mouse sperm parameters..... | 44 |
| 3.1.3 Effects of Aroclor 1254 on mouse epididymal histology | 46 |
| 3.1.4 Endocrine disruptin effets on Aroclor 1254 on mouse epididymis | 49 |
| 3.2 Effects of Aroclor 1254 on gene expression profiles in mouse caput epididymis..... | 51 |
| 3.2.1 Differential expressed genes inuced by different dose of Aroclor 1254 exposure | 51 |
| 3.2.2 PCR validation of differential expressed genes | 53 |
| 3.2.3 Functional annotation of differential expressed genes..... | 54 |
| 3.3 Effects of Aroclor 1254 on function and regulation of epididymal tight junction | 61 |

| | |
|--|-----------|
| 3.3.1 Effects of Aroclor 1254 on expression of epididymal small GTP binding proteins | 61 |
| 3.3.2 Effects of Aroclor 1254 on epididymal actin cytoskeleton | 62 |
| 3.3.3 Effects of Aroclor 1254 on epididymal tight junction | 64 |
| 3.4 Screen of biomarker candidates of PCB exposure..... | 68 |
| 3.4.1 Biomarker candidates from differential expressed genes | 68 |
| 3.4.2 Dose effects of Aroclor 1254 exposure on the Psap expression..... | 69 |
| 3.4.3 Correlation between Psap expression, sperm parameters and PCB accumulation | 71 |
| 3.4.4 Temporal effects of Aroclor 1254 exposure on the Psap expression.... | 72 |
| Chapter 4 Discussion | 73 |
| 4.1 Accumulation of Aroclor 1254 in mouse liver | 73 |
| 4.2 Effects of Aroclor 1254 exposure on body weight and sperm parameters | 73 |
| 4.3 Disruption of Aroclor 1254 exposure on androgen and estrogen function | 74 |
| 4.4 Effects of Aroclor 1254 exposure on gene expression profiles | 76 |
| 4.4.1 Differential expressed genes induced by Aroclor 1254 exposure | 76 |
| 4.4.2 Effects of Aroclor 1254 exposure on gene expression patterns..... | 77 |
| 4.4.3 The comparison of mechanisms in different doses Aroclor 1254 exposure | 78 |
| 4.5 Effects of Aroclor 1254 exposure on actin cytoskeleton | 81 |
| 4.6 Effects of Aroclor 1254 exposure on tight junction..... | 84 |
| 4.7 Screen of biomarker candidate of PCB reproductive toxicology | 88 |
| Chapter 5 Conclusion and prospect..... | 91 |
| 5.1 Conclusion | 91 |
| 5.2 Innovation..... | 92 |
| 5.3 Prospect..... | 92 |
| References | 94 |

摘 要

多氯联苯 (Polychlorinated biphenyls, PCBs) 是最难以降解的持久性有机污染物之一。有关 PCB 的生殖毒性效应及机制已开展了较多的研究, 但 PCB 对附睾的毒性效应及机制还不清楚。本研究以 C57BL/6 品系青春期雄性小鼠为研究对象, 以商品化的 Aroclor 1254 为 PCB 代表, 研究环境水平的 PCB 暴露 (0.5、5、50 和 500 $\mu\text{g}/\text{kg}$) 对附睾的毒性效应及机制。

青春期雄性小鼠在 Aroclor 1254 暴露 50 天后, 其体重、睾丸和附睾重量没有显著性变化, 精子密度、精子活力以及顶体酶活性呈现剂量依赖性下降, 精子畸形率呈现剂量依赖性上升。血清中的雌激素水平表现出剂量依赖性的下降, 雄激素的水平没有表现出显著性的变化。Western blot 检测的结果显示: 附睾中的雌激素受体 α (Estrogen receptor α , ER α) 和雌激素受体 β (Estrogen receptor β , ER β) 并无显著性变化, 而雄激素受体在 500 $\mu\text{g}/\text{kg}$ 组表现出显著性下降, 提示 Aroclor 1254 对附睾表现出抗雄激素效应。

利用 cDNA 微阵列 (Microarray) 技术研究附睾头部在低浓度 (5 $\mu\text{g}/\text{kg}$) 和高浓度 (500 $\mu\text{g}/\text{kg}$) Aroclor 1254 暴露下基因的差异表达, 结果显示: 共筛选出 2350 个差异表达基因, 其中低浓度组引起附睾头部 572 个基因的差异表达, 包括上调基因 212 个, 下调基因 360 个; 高浓度组引起附睾头部 1769 个基因的差异表达, 其中上调基因 433 个, 下调基因 1336 个。应用实时荧光定量 PCR (Real-time fluorescent quantitative PCR, Real-time PCR) 对部分差异表达基因进行了 mRNA 水平的验证, 结果证明了芯片检测的可靠性。进一步应用分子注释系统 (Molecule Annotation System, MAS), 结合功能注释聚类 (Functional Annotation Clustering) 以及聚类分析 (Clustering) 等生物信息技术, 对差异表达基因信息和功能以及所参与的信号通路进行挖掘和注释, 得到差异基因中最为显著的功能类别和信号通路, 其中涵盖了 GTP 结合、核小体组装、能量代谢、RNA 加工等方面。为进一步探索 PCB 对附睾的毒性效应机制提供了依据。

根据功能注释聚类的结果, 发现与 GTP 结合相关的基因在高浓度 Aroclor 1254 暴露后表现出最高的富集度 (Enrichment score), 表明相关基因的表达变化

是 Aroclor 1254 暴露作用于附睾表达谱的最显著的效应之一。MAS 分析表明相关基因的功能包括了紧密连接和肌动蛋白细胞骨架的调控。Western blot 检测表明相关的小 GTP 结合蛋白, 如 K-ras、Cdc42 以及 RhoA 等的表达水平在 PCB 暴露后表现出显著性的下调。小 GTP 结合蛋白通过调节肌球蛋白调节轻链(Myosin regulatory light chain, MRLC) 和丝切蛋白(Cofilin) 的磷酸化水平来调节紧张纤维的收缩和肌动蛋白聚合的稳定性。Western blot 结果表明 MRLC 磷酸化水平上升而 Cofilin 磷酸化水平下降, 提示紧张纤维的收缩加强、肌动蛋白聚合的稳定性增加。

小 GTP 结合蛋白表达的变化和纤维的收缩加强提示紧密连接功能的改变。Western blot 结果表明紧密连接的组成蛋白(胞质紧密粘连蛋白 1) ZO-1 在 500 $\mu\text{g/kg}$ PCB 暴露下表现出显著性的下调。免疫组化结果表明定位与细胞顶端连接的 ZO-1 染色并没有明显的变化。免疫共沉淀的结果表明, Occludin 的酪氨酸磷酸化的水平上升。电镜下的硝酸镧示踪提示紧密连接的透过性增加。紧密连接的透过性关系到精子成熟环境的维持, PCB 暴露对紧密连接的影响提示了 PCB 对生殖毒性的新机制。

对低浓度和高浓度都能够显著性诱导的差异基因进行基因分类学注释(Gene Ontology, GO), 选取 GO 注释包括细胞外空间(Extracellular space 或 Extracellular region) 为候选标志物。从中选取鞘脂激活蛋白原(Prosaposin, Psap) 进行蛋白表达水平的验证。将 Psap 表达水平与精子指标和肝脏中的 PCB 浓度建立相关性, 结果表明 Psap 表达与 PCB 浓度以及精子畸形率负相关而与精子活力正相关。提示 Psap 可能作为 PCB 暴露和雄性生殖毒性的候选标志物。

关键词: 多氯联苯, 附睾, 生殖毒性

ABSTRACT

Polychlorinated biphenyls (PCBs) are among the most persistent environmental pollutants and the reproductive toxicity of PCBs has been extensively studied. However, the toxicological effects and mechanisms of PCBs on epididymis remain unclear. In the present study, toxicological effects and mechanisms of PCB exposures of environmental levels on epididymis of pubertal C57BL/6 mice were investigated.

Male pubertal C57BL/6 mice were exposed to Aroclor 1254, a commercial PCB mixture once widely used by gavage at low dose (0.5, 5, 50 and 500 $\mu\text{g/kg/3days}$, respectively). After exposure for 50 days, the epididymal sperm count, sperm motility and acrosin activity showed a dose-dependent decrease, and the sperm abnormality showed a dose-dependent increase, while no changes in body weight, testis weight or epididymis weight were observed. Serum estrogen level decreased in a dose-dependent manner and the alteration of serum androgen level remained insignificant. The results of Western blot analysis demonstrated that the expression of androgen receptor (AR) in caput epididymis were significantly decreased in 500 $\mu\text{g/kg}$ group while the expression of estrogen receptor α (ER α) and estrogen receptor β (ER β) remained unchanged. It suggested an antiandrogen effect of Aroclor 1254 on epididymis in low-dose PCB exposure.

cDNA microarray technique was used to investigate the gene expression profiles of caput epididymis in response to low dose (5 $\mu\text{g/kg}$) and high dose (500 $\mu\text{g/kg}$) Aroclor 1254 exposure and 2350 differential expressed genes identified. After exposed to low dose Aroclor 1254, 572 genes were differentially regulated, which included 212 up-regulated genes and 360 down-regulated genes. After exposed to high dose Aroclor 1254, 1769 genes were differentially regulated, which included 433 up-regulated genes and 1336 down-regulated genes. Real-time fluorescent quantitative PCR (qPCR) were performed on selected genes for the confirmation of microarray data. The results were in agreement with the microarray data.

Bioinformatics tools including functional annotation clustering and molecule annotation system (MAS) were applied to reveal the biological significance of the gene expression data. The differential expressed genes were involved in a variety of function categories and biological pathways, such as GTP binding, nucleosome assembly, energy metabolism, RNA processing etc.

The results of functional annotation clustering indicated that the functional category related to GTP binding showed the highest enrichment score among the functional categories affected by 500 µg/kg Aroclor 1254 exposure. It suggested that changes in the expression of genes related to GTP binding are among the most significant effects of Aroclor 1254 on epididymis. MAS analysis indicated that GTP binding proteins may involved in many important biological processes, such as regulation of actin cytoskeleton and assembly of tight junction. Western blot analysis revealed significant decreases in protein abundance of small GTP binding proteins: Cdc42, Ras and RhoA after PCB exposure. Alteration in these small GTP binding proteins may affect the contraction of stress fiber and stability of actin filament by affecting the phosphorylation status of their down stream effectors: myosin regulatory light chain (MLC) and cofilin. The increase of phosphorylated MLC and decrease of phosphorylated cofilin were detected by western blot. These results suggested an increase in stress fiber contraction and a decrease in actin filament stability.

The GTP binding protein related actin cytoskeleton alteration may suggest a disruption on tight junction function. Western blot indicated a significant decrease in ZO-1 expression and a trend of decrease in occludin expression. Immunohistochemistry showed a disruption of apical occludin expression and no observable changes in apical ZO-1 expression. Immunoprecipitation assay showed an increase in tyrosine phosphorylated occludin. Lanthanum nitrate tracing under the electron microscope revealed an increased permeability of apical tight junction after 500 µg/kg Aroclor 1254 treatment. The apical tight junction permeability in epididymal principal cells is important to the maintenance of microenvironment which supports the sperm maturation. The disruption on epididymal tight junction may be involved in a novel mechanism by which PCBs affect the post-testicular

sperm maturation.

Gene ontology (GO) annotations were performed on differential expressed genes which were both regulated by 5 µg/kg exposure and 500 µg/kg exposure. The genes related to the GO term: extracellular space or extracellular region were selected as biomarker candidates. Among these candidates, the protein abundance of prosaposin (psap) was examined in epididymis and sperm by western blot and showed a dose-dependent decrease. The expression level of psap in sperm was negatively correlated to hepatic PCB accumulation and sperm abnormality and was positively correlated to sperm motility. The results suggested that psap is potential biomarker of PCB male reproductive toxicity.

Key words: Polychlorinated biphenyls; Epididymis; Reproductive toxicity

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库